

**South African National Essential Medicine List
Adult Hospital Level Medication Review Process
Component: HIV and AIDs**

MEDICINE REVIEW

Executive Summary

Date: 06 February 2020
Medicine (INN): Tenofovir alafenamide (TAF)
Medicine (ATC): J05AF13
Indication (ICD10 code): B20
Patient population: HIV-1 infected adult patients
Prevalence of condition: An estimated 7.02 million people were living with HIV in South Africa in 2016, representing 12.7% of the national population or 19.1% of those aged 15-49 years(1)
Level of Care: Primary level of care
Prescriber Level: Nurse prescriber, doctor
Current standard of Care: Tenofovir disoproxil fumarate (TDF)
Efficacy estimates: At 48 weeks, TAF was non-inferior to TDF, viral suppression rates (HIV RNA <50 copies/mL) were 92% vs. 90% respectively.
 Numbers needed to treat** for TAF vs. TDF, based on VL suppression estimates from the RCT by Sax et al, 2015: NNT = 50 (95% CI 21 - 142) (About 2% absolute risk reduction (2))
 **NNT = 1/Absolute RD.
Motivator/reviewer name(s): Dr S Takuva, Mr NJ Nabyoma, Prof G Maartens
PTC affiliation: N/A

Name of author(s)/motivator(s): Dr S Takuva, Mr NJ Nabyoma, Prof G Maartens

Authors affiliation and conflict of interest details:

Primary reviewers:

Dr S Takuva: No applicable conflict of interest to declare

- 1) School of Health Systems and Public Health, Faculty of Health Sciences, University of Pretoria, South Africa
- 2) Perinatal HIV Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
- 3) Adult Hospital Level Committee, 2017-2020

Mr NJ Nabyoma: No applicable conflict of interest to declare

- 1) Department of Health, North West Province, South Africa.
- 2) Adult Hospital Level Committee, 2017-2020

Secondary reviewer:

Prof G Maartens: No applicable conflict of interest to declare

- 1) Department of Pharmacology, University of Cape Town, South Africa
- 2) National Essential Medicines List Committee, 2017-2020

Introduction

Since April 2010, Tenofovir disoproxil fumarate (TDF) has been the mainstay of first line antiretroviral treatment (ART) in South Africa.(3) It is generally well-tolerated, however, long-term use of TDF is associated with progressive declines in glomerular function and chronic kidney disease in HIV-infected patients.(4–11) Data from a large ART cohort in South Africa showed that patients with mild or moderate renal dysfunction were at higher risk of nephrotoxicity, while those with mild or moderate renal dysfunction vs. normal renal function were at highest risk of death by 48-months of follow-up.(5) In another South African cohort study with over 15,000 patients on TDF containing regimens followed up for a median duration of 13 months, patients without

renal impairment at baseline (eGFR ≥ 90 mL/min) experienced small but significant declines in eGFR over time.(12) In another study from 1092 HIV-infected patients initiating tenofovir at a primary care clinic in Cape Town, South Africa, renal function was assessed for the first 12 months on ART, generally, renal function improved in the study population during the first year on ART. Renal impairment during the first 12 months of tenofovir-containing ART was 3%.(11) However, the burden of chronic kidney disease among HIV-infected patients in South Africa is high (6%) and estimates indicate that approximately 10% of patients (an estimated 702,000 patients from current HIV prevalence figures) will suffer from HIV-related renal failure or renal toxicities throughout the course of their disease.(5)(13)(14)

Whilst data on the prevalence and sequelae of metabolic bone diseases among HIV-infected patients in resource-limited settings like South Africa is scanty(15), a meta-analysis reported a 60% increased fracture risk in HIV-infected individuals when compared to uninfected individuals.(16) Patients treated with TDF have been observed to have greater decline in bone mineral density (BMD) relative to some other NRTIs.(16–21)

Tenofovir alafenamide (TAF), an oral prodrug of tenofovir, is now included as a component of several recommended first-line antiretroviral therapy regimens. These recommendations are based on data from comparative trials demonstrating that TAF-containing regimens are as effective in achieving or maintaining virologic suppression as TDF-containing regimens but with more favourable effects on markers of renal and bone health.(2,22–29) Unlike TDF, which should be avoided or dose-adjusted in patients with renal dysfunction or estimated creatinine clearance (CrCl) < 80 mL/min, TAF-containing regimens appear to be safe and are FDA approved for use in patients with estimated CrCl as low as 30 mL/min.

The aim of this medicine review is to review current available evidence for the use of TAF as part of first line antiretroviral therapy in a roll-out antiretroviral therapy programme.

Question:

1. TAF is non-inferior to TDF as part of ART regimen to treat HIV-1 infection
2. TAF has a better safety profile to TDF (especially renal and bone)

PICOT criteria;

Population	HIV-1 infected adult patients
Intervention	Tenofovir alafenamide
Comparison	Tenofovir disoproxil fumarate either as comparison arm or switch study
Outcomes	Mortality, AIDS progression, Viral suppression, Immunological response, Adverse events and severity,

Search strategy:

An electronic literature search of the PubMed and EMBASE database from beginning of time till 30 January 2020 was undertaken using different combinations of:

("HIV"[MeSH Terms] OR "HIV"[All Fields]) AND ("tenofovir disoproxil fumarate"[All Fields] OR TDF [All Fields])) AND ("tenofovir alafenamide"[All Fields] OR TAF [All Fields])

WHO HIV treatment guidelines also reviewed, as they are relevant to this setting.

Selection of Studies:

Abstracts from 180 publications were screened.

Exclusions were;

- Out of 29 review articles, 15 were excluded – did not compare TAF to TDF
- Out of 69 publications, 57 excluded as they were not randomized clinical trials or systematic reviews
- To avoid repetition, review articles (including systematic reviews) were scanned if they included identified RCTs)

Four meta-analyses and an expert think tank review commissioned by the WHO were finally selected for evidence synthesis.

Evidence Synthesis:

The efficacy and safety of TAF-containing regimens vs. TDF-containing regimens have been mostly evaluated in the context of the coformulation of elvitegravir, cobicistat, emtricitabine and darunavir. Comprehensive reviews were identified that included RCTs published to date of synthesis. While there is some overlap of studies in these systematic reviews below, this is little as some reviews focussed on switch studies and others focussed on direct parallel TDF vs. TAF comparisons. Where a review mainly updated a previously published review, the previous one was excluded to reduce duplication.

Systematic Reviews and Review Articles:

Tao et al 2020: Seven RCTs with a total of 6269 participants.

- Virologic suppression rates were similar: (RR, 1.02; 95% CI, 1.00-1.04; $p > 0.05$) at week 24 (94.0% vs. 94.2%), week 48 (90.7% vs. 89.5%), and week 96 (86.2% vs. 84.8%).
- Both treatments were safe and well-tolerated, and most adverse events were similar as mild to moderate in severity.
- Compared with the TDF-containing regimens, the TAF-containing regimens in patients had significantly smaller reductions in both hip (RR, 0.33; 95CI, 0.29-0.39; $p < 0.05$) and spine (RR, 0.58; 95CI, 0.51-0.65; $p < 0.05$).
- Additionally, the TAF-containing regimens in patients had significantly fewer increases for renal events than those of the TDF-containing regimens through 48 weeks (0.31; 95% CI, 0.18-0.55; $p < 0.05$).

Tao et al 2019: Eight eligible phase III RCTs included and altogether 7613 patients were recruited.

- patients switched to TAF-containing regimens had significantly better viral suppression than those continuing TDF-containing regimens at weeks 48 and 96 (RR, 1.02; 95CI, 1.00-1.03), but no significant difference in the per-protocol (PP) analysis (RR, 1.00; 95CI, 0.99-1.01).
- Compared with those receiving the TDF-containing regimens, virologically suppressed HIV-infected patients on the TAF-containing regimens had significant increases in CD4 cell counts (SMD, 0.12; 95CI, 0.08 to 0.17), renal and bone parameters at the hip (RR, 2.86; 95CI, 2.24-3.64) and the spine (RR, 2.43; 95 CI, 2.03-2.90) between weeks 48 and 96.
- Among these RCTs, 5.2% of all participants in the TAF-containing regimens and 3.8% of all participants in the TDF-containing regimens started lipid-lowering drugs, and no statistical differences were found between the two groups after 48 weeks and 96 weeks of treatment (RR, 1.27; 95%CI, 0.94–1.71)

Tamuzi et al 2018: 18 randomized controlled trials were used in the Meta-analysis and these are the finding

- HIV-infected patients on TAF based regimens reduced HIV-RNA<50RNAc/ml by 13% compared to TDF contained group ($P=0.02$)
- TAF to TDF based regimens, the glomerular filtration rate yielded a pooled MD estimate of -3.94(-6.07 to -1.81, $P<0.000001$)
- The MD of percentage change hip bone mineral density was decreased in TDF compared to TAF -1.93 with $P<0.00001$. MD of percentage change spine bone mineral density was decreased in TDF compared to TAF -1.77 (-1.97 to -1.58) with $P=0.001$.
- Adverse events and serious adverse events were not significant in both TAF and TDF groups.

Gotham et al 2017: The authors identified 10 randomized controlled trials comparing TDF with TAF (6969 patients, 8043 patient-years of follow-up). (23)The key points from this Meta-analysis were;

- There were no significant differences in treatment efficacy, resistance, or adverse events between TAF and TDF arms.
- There were significant differences, favouring TAF, in BMD and renal function measures, but no significant differences in treatment discontinuations because of bone or renal toxicity.
- TAF treatment was associated with higher HDL levels. A few patients were started on statins.
- There is a lack of data for safety of TAF in pregnancy, TB co-infection, and patients with low CD4 count (<50 cells/mm3).

Vitoria M et al 2017: There were 60 experts invited, including members of the WHO HIV Guidelines committee, specialists in paediatrics and HIV drug resistance, UNITAID, the Clinton Health Access Initiative, USAID, Centres for Disease Control and PEPFAR. The two main questions discussed at this WHO Think-Tank meeting were;

- Is there already enough evidence to support the efficacy and safety of DTG, TAF and EFV400 to justify their use in millions of people in LMICs?
- What clinical trials and pharmacovigilance studies are needed to assess drug safety when these new treatments are used more widely.(31)

These were the key points from the think tank;

- It was agreed that additional safety and efficacy data on DTG, TAF and EFV400 in some subpopulations are needed, particularly for pregnant women and people with HIV–TB coinfection.
- At the meeting, there was limited support for the introduction of TAF as part of first-line antiretroviral treatment in low-income and middle-income settings.
- There was an overall agreement for 6-monthly reviews of safety and efficacy data, in parallel with a phased introduction of the new antiretrovirals.

EVIDENCE TO DECISION FRAMEWORK

	JUDGEMENT	SUPPORTING EVIDENCE & ADDITIONAL CONSIDERATIONS
QUALITY OF EVIDENCE	<p>What is the overall confidence in the evidence of effectiveness?</p> <p>Confident Not confident Uncertain</p> <p><input type="checkbox"/> <input type="checkbox"/> <input checked="" type="checkbox"/></p>	<p>Rationale (also see under Recommendations):</p> <p>-The efficacy and safety of TAF-containing regimens vs. TDF-containing regimens have been mostly evaluated in the context of the coformulation of elvitegravir, cobicistat, emtricitabine and darunavir. There is insufficient data where it has been evaluated in standard formulation used in LMICs.</p> <p>-The synthesis below shows that TAF is no more effective than TDF. TAF overall, shows slightly lower toxicity in these studies especially regarding renal and bone health markers – the clinical significance of these differences in markers is not clear. However, these findings should be interpreted cautiously as in most studies TAF was co-formulated with cobicistat, where the TAF dose is reduced from 25mg to 10mg. There is a need for trials comparing or evaluating efficacy and especially safety of TAF head for head in standard coformulations used in low middle-income countries.</p> <p>-Emerging observational data suggests switching from TDF to TAF may cause a statistically significant worsening of the lipid profile that may have clinical relevance. This is likely seen in patients with cardiovascular risk factors such as older age and high BMI. The lower concentrations of TDF in plasma from TAF as compared with TDF, and the lipid-lowering effect of TDF may explain the increases in total cholesterol in the TAF group compared with the TDF group. It may be important to weigh the possible benefit of lipid changes associated with TDF against the possible benefit of TAF for bone and kidney.</p>
BENEFITS & HARMS	<p>Do the desirable effects outweigh the undesirable effects?</p> <p>Benefits outweigh harms Harms outweigh benefits Benefits = harms or Uncertain</p> <p><input type="checkbox"/> <input type="checkbox"/> <input checked="" type="checkbox"/></p>	<p>See above.</p>

THERAPEUTIC INTERCHANGE	Therapeutic alternatives available: Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> List the members of the group. Other NRTIs like TDF, ABC List specific exclusion from the group: n/a	Rationale for therapeutic alternatives included: Other NRTIs References: n/a					
	Is there important uncertainty or variability about how much people value the options? Minor <input type="checkbox"/> Major <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/> Is the option acceptable to key stakeholders? Yes <input type="checkbox"/> No <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/>						
RESOURCE USE	How large are the resource requirements? More intensive <input type="checkbox"/> Less intensive <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/>	Cost of medicines/ month: <table border="1"> <thead> <tr> <th>Medicine</th> <th>Cost (ZAR)</th> </tr> </thead> <tbody> <tr> <td>Not currently SAHPRA registered.</td> <td></td> </tr> </tbody> </table> Additional resources: n/a		Medicine	Cost (ZAR)	Not currently SAHPRA registered.	
Medicine	Cost (ZAR)						
Not currently SAHPRA registered.							
EQUITY	Would there be an impact on health inequity? Yes <input type="checkbox"/> No <input type="checkbox"/> Uncertain <input checked="" type="checkbox"/>						
FEASIBILITY	Is the implementation of this recommendation feasible? Yes <input checked="" type="checkbox"/> No <input type="checkbox"/> Uncertain <input type="checkbox"/>						

Type of recommendation	We recommend against the option and for the alternative	We suggest not to use the option or to use the alternative	We suggest using either the option or the alternative	We suggest using the option	We recommend the option
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Recommendation: TAF not be considered for inclusion in the Adult Hospital Level EML, currently.

Note:

- Based on the best available evidence, TAF is no better in efficacy than TDF and may have small safety benefits whose clinical relevance is still uncertain. TAF can be considered in first line regimens in the future should the TAF/FTC co-formulation or FDCs be licensed in RSA (FTC/TAF/DTG) – for patients with contraindications to TDF i.e. advanced renal disease.
- There is very limited clinical experience of TAF in pregnancy and we therefore do not recommend TAF use in pregnancy.
- The potential for the interaction of TAF with rifampicin exists and concurrent therapy still needs further evaluation.

Rationale:

- The efficacy and safety of TAF-containing regimens vs. TDF-containing regimens have been mostly evaluated in the context of the coformulation of elvitegravir, cobicistat, emtricitabine and darunavir. There is insufficient data where it has been evaluated in standard formulation used in LMICs.
- The synthesis shows that TAF is no more effective than TDF. TAF overall, shows slightly lower toxicity in these studies especially with regard to renal and bone health markers – the clinical significance of these differences in markers is not clear. However, these findings should be interpreted cautiously as in most studies TAF was co-formulated with cobicistat, where the TAF dose is reduced from 25mg to 10mg. There is a need for trials comparing or evaluating efficacy and especially safety of TAF head for head in standard coformulations used in low middle-income countries.
- Emerging observational data suggests switching from TDF to TAF may cause a statistically significant worsening of the lipid profile that may have clinical relevance. This is likely seen in patients with cardiovascular risk factors such as older age and high BMI. The lower concentrations of TDF in plasma from TAF as compared with TDF, and the lipid-lowering effect of TDF may explain the increases in total cholesterol in the TAF group compared with the TDF group. It may be important to weigh the possible benefit of lipid changes associated with TDF against the possible benefit of TAF for bone and kidney.

Level of Evidence: I Systematic Reviews and Meta-Analysis of Randomized Clinical Trials

NEMLC MEETING OF 19 MARCH 2019:

NEMLC accepted this evidence review and the proposal as recommended by the Adult Hospital Level Expert Review Committee, above.

NEMLC also acknowledged that TAF-containing fixed-dose combination formulations are currently not SAHPRA registered and thus not currently available on the South African market. The current antiretroviral recommendations, as recommended in the Standard Treatment Guidelines (Adult Hospital Level, 2019 edition) and National HIV Guidelines, 2019 edition are sufficient.

Proposed TAF-containing antiretroviral regimens - refer to Annexure A.

Review indicator:

Evidence of efficacy	Evidence of harm	Price reduction
<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

VEN status: n/a

Vital	Essential	Necessary
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Monitoring and evaluation considerations: n/a

Research priorities

Safety and efficacy in pregnancy and HIV-TB co-treatment.
Long term safety data using coformulation used in LMICs

References

1. Johnson LF, Chiu C, Myer L, Davies M-A, Dorrington RE, Bekker L-G, et al. Prospects for HIV control in South Africa: a model-based analysis. *Glob Health Action*. 2016 Jan;9(1):30314.
2. Sax PE, Wohl D, Yin MT, Post F, DeJesus E, Saag M, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate, coformulated with elvitegravir, cobicistat, and emtricitabine, for initial treatment of HIV-1 infection: two randomised, double-blind, phase 3, non-inferiority trials. *Lancet* (London, England). 2015 Jun;385(9987):2606–15.
3. National Department of Health. The South African Antiretroviral Treatment Guidelines [Internet]. 2010. Available from: <http://www.doh.gov.za/docs/factsheets/guidelines/art.pdf>
4. Franey C, Knott D, Barnighausen T, Dedicoat M, Adam A, Lessells RJ, et al. Renal impairment in a rural African antiretroviral programme. *BMC Infect Dis*. 2009 Aug;9:143.
5. Brennan A, Evans D, Maskew M, Naicker S, Ive P, Sanne I, et al. Relationship between renal dysfunction, nephrotoxicity and death among HIV adults on tenofovir. *AIDS*. 2011 Aug;25(13):1603–9.
6. Seedat F, Martinson N, Motlhaoleng K, Abraham P, Mancama D, Naicker S, et al. Acute Kidney Injury, Risk Factors, and Prognosis in Hospitalized HIV-Infected Adults in South Africa, Compared by Tenofovir Exposure. *AIDS Res Hum Retroviruses*. 2017 Jan;33(1):33–40.
7. Le MP, Landman R, Koulla-Shiro S, Charpentier C, Sow P-S, Diallo M-B, et al. Tenofovir plasma concentrations related to estimated glomerular filtration rate changes in first-line regimens in African HIV-infected patients: ANRS 12115 DAYANA substudy. *J Antimicrob Chemother*. 2015 May;70(5):1517–21.
8. Nelson MR, Katlama C, Montaner JS, Cooper DA, Gazzard B, Clotet B, et al. The safety of tenofovir disoproxil fumarate for the treatment of HIV infection in adults: the first 4 years. *AIDS*. 2007 Jun;21(10):1273–81.
9. Cooper RD, Wiebe N, Smith N, Keiser P, Naicker S, Tonelli M. Systematic review and meta-analysis: renal safety of tenofovir disoproxil fumarate in HIV-infected patients. *Clin Infect Dis*. 2010 Sep;51(5):496–505.
10. Woodward CLN, Hall AM, Williams IG, Madge S, Copas A, Nair D, et al. Tenofovir-associated renal and bone toxicity. *HIV Med*. 2009 Sep;10(8):482–7.
11. Kamkuemah M, Kaplan R, Bekker L-G, Little F, Myer L. Renal impairment in HIV-infected patients initiating tenofovir-containing antiretroviral therapy regimens in a Primary Healthcare Setting in South Africa. *Trop Med Int Health*. 2015 Apr;20(4):518–26.
12. De Waal R, Cohen K, Fox MP, Stinson K, Maartens G, Boule A, et al. Changes in estimated glomerular filtration rate over time in South African HIV-1-infected patients receiving tenofovir: a retrospective cohort study. *J Int AIDS Soc*. 2017 Apr;20(1):1–8.
13. G B. HIV-related renal disease: a clinical and practical approach in the South African context. *South Africa Med J*. 2003;(Feb):11–4.
14. Franceschini N, Napravnik S, Eron JJJ, Szczech LA, Finn WF. Incidence and etiology of acute renal failure among ambulatory HIV-infected patients. *Kidney Int*. 2005 Apr;67(4):1526–31.
15. Matovu FK, Watanachanya L, Beksinska M, Pettifor JM, Ruxrungtham K. Bone health and HIV in resource-limited settings: a scoping review. *Curr Opin HIV AIDS*. 2016 May;11(3):306–25.
16. Shiao S, Broun EC, Arpadi SM, Yin MT. Incident fractures in HIV-infected individuals: a systematic review and meta-analysis. *AIDS*. 2013 Jul;27(12):1949–57.
17. Stellbrink H-J, Orkin C, Arribas JR, Compston J, Gerstoft J, Van Wijngaerden E, et al. Comparison of changes in bone density and turnover with abacavir-lamivudine versus tenofovir-emtricitabine in HIV-infected adults: 48-week results from the ASSERT study. *Clin Infect Dis*. 2010 Oct;51(8):963–72.
18. Grant PM, Cotter AG. Tenofovir and bone health. *Curr Opin HIV AIDS*. 2016 May;11(3):326–32.
19. Bedimo RJ, Drechsler H, Jain M, Cutrell J, Zhang S, Li X, et al. The RADAR study: week 48 safety and efficacy of Raltegravir combined with boosted DARunavir compared to tenofovir/emtricitabine combined with boosted darunavir in antiretroviral-naïve patients. Impact on bone health. *PLoS One*. 2014;9(8):e106221.
20. Gallant JE, Staszewski S, Pozniak AL, DeJesus E, Suleiman JMAH, Miller MD, et al. Efficacy and safety of tenofovir DF vs stavudine in combination therapy in antiretroviral-naïve patients: a 3-year randomized trial. *JAMA*. 2004 Jul;292(2):191–201.
21. McComsey GA, Kitch D, Daar ES, Tierney C, Jahed NC, Tebas P, et al. Bone mineral density and fractures in antiretroviral-naïve persons randomized to receive abacavir-lamivudine or tenofovir disoproxil fumarate-emtricitabine along with efavirenz or atazanavir-ritonavir: Aids Clinical Trials Group A5224s, a substudy of ACT. *J Infect Dis*. 2011 Jun;203(12):1791–801.
22. Orkin C, DeJesus E, Ramgopal M, Crofoot G, Ruane P, LaMarca A, et al. Switching from tenofovir disoproxil fumarate to tenofovir alafenamide coformulated with rilpivirine and emtricitabine in virally suppressed adults with HIV-1 infection: a randomised, double-blind, multicentre, phase 3b, non-inferiority study. *Lancet HIV*. 2017 May;4(5):e195–204.
23. Gotham D, Hill A, Pozniak AL. Candidates for inclusion in a universal antiretroviral regimen: tenofovir alafenamide. *Curr Opin HIV AIDS*. 2017 Apr;
24. Sax PE, Zolopa A, Brar I, Elion R, Ortiz R, Post F, et al. Tenofovir alafenamide vs. tenofovir disoproxil fumarate in single tablet regimens for initial HIV-1 therapy: a randomized phase 2 study. *J Acquir Immune Defic Syndr*. 2014 Sep;67(1):52–8.
25. Gallant JE, Daar ES, Raffi F, Brinson C, Ruane P, DeJesus E, et al. Efficacy and safety of tenofovir alafenamide versus tenofovir disoproxil fumarate given as fixed-dose combinations containing emtricitabine as backbones for treatment of HIV-1 infection in virologically suppressed adults: a randomised, double-blind, active. *Lancet HIV*. 2016 Apr;3(4):e158–65.
26. Arribas JR, Thompson M, Sax PE, Haas B, McDonald C, Wohl DA, et al. Brief Report: Randomized, Double-Blind Comparison of Tenofovir Alafenamide (TAF) vs Tenofovir Disoproxil Fumarate (TDF), Each Coformulated With Elvitegravir, Cobicistat, and Emtricitabine (E/C/F) for Initial HIV-1 Treatment: Week 144 Results. *J Acquir Immune Defic Syndr*. 2017 Jun;75(2):211–8.
27. Wohl D, Oka S, Clumeck N, Clarke A, Brinson C, Stephens J, et al. Brief Report: A Randomized, Double-Blind Comparison of Tenofovir Alafenamide Versus Tenofovir Disoproxil Fumarate, Each Coformulated With Elvitegravir, Cobicistat, and Emtricitabine for Initial HIV-1 Treatment: Week 96 Results. *J Acquir Immune Defic Syndr*. 2016 May;72(1):58–64.
28. Mills A, Arribas JR, Andrade-Villanueva J, DiPerri G, Van Lunzen J, Koenig E, et al. Switching from tenofovir disoproxil fumarate to tenofovir alafenamide in antiretroviral regimens for virologically suppressed adults with HIV-1 infection: a randomised, active-controlled, multicentre, open-label, phase 3, non-inferiority study. *Lancet Infect Dis*. 2016 Jan;16(1):43–52.

29. Pozniak A, Arribas JR, Gathe J, Gupta SK, Post FA, Bloch M, et al. Switching to Tenofovir Alafenamide, Coformulated With Elvitegravir, Cobicistat, and Emtricitabine, in HIV-Infected Patients With Renal Impairment: 48-Week Results From a Single-Arm, Multicenter, Open-Label Phase 3 Study. *J Acquir Immune Defic Syndr*. 2016 Apr;71(5):530–7.
30. Post FA, Yazdanpanah Y, Schembri G, Lazzarin A, Reynes J, Maggiolo F, et al. Efficacy and safety of emtricitabine/tenofovir alafenamide (FTC/TAF) vs. emtricitabine/tenofovir disoproxil fumarate (FTC/TDF) as a backbone for treatment of HIV-1 infection in virologically suppressed adults: subgroup analysis by third agent of a randomi. *HIV Clin Trials*. 2017 Mar;1–6.
31. Tamuzi JL, Tshimwanga JL, Bulabula AN, Muyaya LM. Tenofovir Alafenamide versus Tenofovir Disoproxil Fumarate: Systematic Review and Meta-Analysis. *Int J Pul & Res Sci*. 2018;2:555600.
32. Vitoria M, Ford N, Clayden P, Pozniak AL, Hill AM. When could new antiretrovirals be recommended for national treatment programmes in low-income and middle-income countries: results of a WHO Think Tank. *Curr Opin HIV AIDS*. 2017 Apr;
33. Gonciulea A, Wang R, Althoff KN, Palella FJ, Lake J, Kingsley LA, et al. An increased rate of fracture occurs a decade earlier in HIV R compared with HIV S men. 2017;(December 2016):1435–43.
33. Venter WDF, Moorhouse M, Sokhela S, Fairlie L, Mashabane N, Masenya M, Serenata C, Akpomiemie G, Qavi A, Chandiwana N, Norris S, Chersich M, Clayden P, Abrams E, Arulappan N, Vos A, McCann K, Simmons B, Hill A. Dolutegravir plus Two Different Prodrugs of Tenofovir to Treat HIV. *N Engl J Med*. 2019 Aug 29;381(9):803-815. doi: 10.1056/NEJMoa1902824. Epub 2019 Jul 24. PubMed PMID: 31339677.
34. Hagins D, Orkin C, Daar ES, Mills A, Brinson C, DeJesus E, Post FA, Morales-Ramirez J, Thompson M, Osiyemi O, Rashbaum B, Stellbrink HJ, Martorell C, Liu H, Liu YP, Porter D, Collins SE, SenGupta D, Das M. Switching to coformulated rilpivirine (RPV), emtricitabine (FTC) and tenofovir alafenamide from either RPV, FTC and tenofovir disoproxil fumarate (TDF) or efavirenz, FTC and TDF: 96-week results from two randomized clinical trials. *HIV Med*. 2018 Nov;19(10):724-733. doi: 10.1111/hiv.12664. Epub 2018 Aug 12. PubMed PMID: 30101539; PubMed Central PMCID: PMC6221083
35. DeJesus E, Haas B, Segal-Maurer S, Ramgopal MN, Mills A, Margot N, Liu YP, Makadzange T, McCallister S. Superior Efficacy and Improved Renal and Bone Safety After Switching from a Tenofovir Disoproxil Fumarate- to a Tenofovir Alafenamide-Based Regimen Through 96 Weeks of Treatment. *AIDS Res Hum Retroviruses*. 2018 Apr;34(4):337-342. doi: 10.1089/AID.2017.0203. Epub 2018 Mar 20. PubMed PMID: 29368537
36. Raffi F, Orkin C, Clarke A, Slama L, Gallant J, Daar E, Henry K, Santana-Bagur J, Stein DK, Bellos N, Scarsella A, Yan M, Abram ME, Cheng A, Rhee MS. Brief Report: Long-Term (96-Week) Efficacy and Safety After Switching From Tenofovir Disoproxil Fumarate to Tenofovir Alafenamide in HIV-Infected, Virologically Suppressed Adults. *J Acquir Immune Defic Syndr*. 2017 Jun 1;75(2):226-231. doi: 10.1097/QAI.0000000000001344. PubMed PMID: 28272164; PubMed Central PMCID: PMC5427981.
37. Tao X, Lu Y, Zhou Y, Zhang L, Chen Y. Efficacy and safety of the regimens containing tenofovir alafenamide versus tenofovir disoproxil fumarate in fixed-dose single-tablet regimens for initial treatment of HIV-1 infection: A meta-analysis of randomized controlled trials. *Int J Infect Dis*. 2020 Jan 24.
38. Wang H, X Lu, X Yang, N Xu. The efficacy and safety of tenofovir alafenamide versus tenofovir disoproxil fumarate in antiretroviral regimens for HIV-1 therapy: Meta-analysis *Medicine*. 2016 Oct; 95(41): e5146.
39. KJ Kauppinen, P Kivelä, J Sutinen. Switching from Tenofovir Disoproxil Fumarate to Tenofovir Alafenamide Significantly Worsens the Lipid Profile in a Real-World Setting. *AIDS Patient Care and STDs*. Dec 2019.500-506.
40. Milinkovic A, Berger F, Arenas-Pinto A, Mauss S. Lipid changes due to tenofovir alafenamide are reversible by switching back to tenofovir disoproxil fumarate. *AIDS*. 2019 Aug 29. doi: 10.1097
41. Cid-Silva P, Fernández-Bargiela N, Margusino-Framiñán L, Balboa-Barreiro V, Mena-De-Cea Á, López-Calvo S, Vázquez-Rodríguez P, Martín-Herranz I, Míguez-Rey E, Poveda E, Castro-Iglesias Á. Treatment with tenofovir alafenamide fumarate worsens the lipid profile of HIV-infected patients versus treatment with tenofovir disoproxil fumarate, each coformulated with elvitegravir, cobicistat, and emtricitabine. *Basic Clin Pharmacol Toxicol*. 2019 Apr;124(4):479-490.
42. Tao X, Lu Y, Zhou Y, Huang Y, Chen Y. Virologically suppressed HIV-infected patients on TDF-containing regimens significantly benefit from switching to TAF-containing regimens: A meta-analysis of randomized controlled trials. *Int J Infect Dis*. 2019 Oct; 87:43-53.

ANNEXURE A

Potential TAF-containing regimens

*The following 4 tenofovir alafenamide-containing FDC tablets are FDA-approved for HIV treatment:

- 1) elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (brand name: Genvoya®),
- 2) emtricitabine/rilpivirine/tenofovir alafenamide (brand name: Odefsey®), and
- 3) Descovy®. As a stand-alone agent, tenofovir alafenamide (brand name: Vemlidy®) is FDA-approved for chronic hepatitis B virus (HBV) infection treatment.
- 4) Dolutegravir/emtricitabine/tenofovir alafenamide (brand name: Kocitaf)

Abbreviations

DTG	Dolutegravir
TDF	Tenofovir disoproxil fumarate
FTC	Emtricitabine
3TC	Lamivudine
ABC	Abacavir
TAF	Tenofovir alafenamide fumarate
EVG/c	Elvitegravir/cobicistat